

Asymmetry in hierarchy model of Bonabeau et al

D. Stauffer¹ and J.S. Sá Martins²

Laboratoire PMMH, Ecole Supérieure de Physique et Chimie Industrielle,
10 rue Vauquelin, F-75231 Paris, Euroland

¹ Visiting from Institute for Theoretical Physics, Cologne University, D-50923 Köln, Euroland; stauffer@thp.uni-koeln.de

² Visiting from Instituto de Física, Universidade Federal Fluminense; Av. Litorânea s/n, Boa Viagem, Niterói 24210-340, RJ, Brazil; jssm@if.uff.br

Abstract: The 1995 model of Bonabeau et al. explains the emergence of social hierarchies through randomness, but gives as many leaders as followers. A simple modification allows a more realistic asymmetry with much less leaders.

Ref. [1] introduced an explanation, henceforth called the Bonabeau model, to explain the emergence of social hierarchies through randomness: If two people compete for the same living space, randomly one of them wins and one of them loses. The more victories (losses) a person had in the recent past, the higher (lower) is the probability to win the duel. With suitable modifications [2, 3] an equilibrium phase transition was found at a concentration near 0.32 between egalitarian society at low population density and hierarchical society at high population density. Schweitzer [4] has criticized that in this model there are as many leaders as there are followers, in contrast to reality. This symmetry is avoided in the present modification.

In the previous version of the Bonabeau model [3], people diffuse randomly on a big square lattice, on which they have a constant density $0 < p < 1$. If one agent i wants to move onto a site already occupied by another agent k , then a fight takes place. It is won with probability q (see below) by agent i , and with probability $1 - q$ by agent k . If k wins, both stay at their place; if k wins the two people exchange their positions. The standard deviation σ of the probabilities is the order parameter:

$$\sigma^2 = \langle q^2 \rangle - \langle q \rangle^2 \quad . \quad (1)$$

Each agent i keeps in mind a history $h(i)$ which is increased by one for every victory and decreased by one for every loss; moreover, the current value of h is diminished by ten percent after every time step. One time step means that on average every agent is selected randomly once to move and possibly

fight. Initially σ and all h are zero. The above probability q for i to win against k is

$$q = 1/(1 + \exp(\sigma[h(k) - h(i)]) \quad . \quad (2)$$

Thus a global feedback exist through the order parameter σ entering Eq. (2).

This model is by definition completely symmetric with respect to victories and losses. We measure the power of each agent on a scale from -20 to $+20$ through the ratio $20h/H$ where H is the maximal value of $|h(i)|$ at that moment in the whole population. Thus we have on average as many people of power -10 as with power $+10$. This unrealistic feature is now avoided by decreasing h after a loss by an amount $F > 1$ while increasing it after a victory only by 1. This factor F thus gives more weight to a loss than to a victory, just as in some football tournaments a loss means the end of participation, while a victory only allows to proceed to the next round.

This asymmetry decreases the phase transition in the concentration p from 0.32 for the old case $F = 1$ to 0.134 at $F = 2$ and 0.006 at $F = 4$, Fig.1. And it produces an asymmetry visible in Fig.2 which already for a factor $F = 2$ gives six times less people with positive power than with negative power. The few people with strongly negative power might be jail inmates. We made 200 or 500 iterations to get equilibrium. Note that the power distribution hardly notices the phase transition, as seen by comparing $+$ with x in Fig. 2. Fig.3 shows that also on the simple cubic lattice a phase transition exists; the power distribution then also looks similar to Fig.2 (not shown).

Acknowledgements: We thank S. Moss de Oliveira for discussion, PMMH at ESPCI for the warm hospitality and Sorin Tănase-Nicola for help with the computer facilities. JSSM acknowledges funding from the Brazilian agencies CNPq and FAPERJ.

References

- [1] E. Bonabeau, G. Theraulaz, J.L. Deneubourg, Physica A 217, 373 (1995)
- [2] A.O. Sousa, D. Stauffer, Int. J. Mod. Phys. C 11, 1063 (2000)
- [3] D. Stauffer, Int. J. Mod. Phys. C 14, 237 (2003)
- [4] F. Schweitzer, priv. comm., September 2002





Figure 1: Order parameter σ versus population density p for factor 2 (part a) and factor 4 (part b).




Figure 2: Histogram of people having a certain power, as measured by $20h/H$. The asymmetry factor is 2 for $p = 0.13$ below the transition (+) and for $p = 0.14$ above the transition (x), while it is 4 for $p = 0.006$ near the transition (stars).




Figure 3: As Fig.1a but in three dimensions.